

Local and global structure of a thick domain wall space-time

*Anzhong Wang*¹

Department of Astrophysics, Observatorio Nacional, Rua General José
Cristino 77, 20921 – 400 Rio de Janeiro, RJ, Brazil

*Patricio S. Letelier*²

Department of Applied Mathematics-IMECC, Universidade Estadual de
Campinas, 13081 – 970 Campinas, SP, Brazil

The local and global properties of the Goetz thick plane domain wall space-time are studied. It is found that when the surface energy of the wall is greater than a critical value σ_c , the space-time will be closed by intermediate singularities at a finite proper distance. A model is presented in which these singularities will give rise to scalar ones when interacting with null fluids. The maximum extension of the space-time of the wall whose surface energy is less than σ_c is presented. It is shown that for certain choice of the free parameter the space-time has a black hole structure but with plane symmetry.

PACS numbers: 04.20.Dw, 04.30.Nk, 97.60.Sm.

¹e-mail address: wang@on.br

²e-mail address: letelier@ime.unicamp.br

The study of topological domain walls is crucial for understanding the inflationary Universe scenario. According to the theory [1], the Universe undertakes an exponential expansion in the early stages of its evolution, triggered by some phase transitions associated with spontaneous symmetry breaking of the Higgs field of some Grand Unified Theory. It was found that such a kind of expansions is needed in order to solve some long-standing puzzles in Cosmology, such as, the horizon, flatness, and monopole. The evolution of the Universe from the exponential expansion to its present Friedmann-Robertson-Walker form is completed by the spontaneous nucleation of bubbles of true vacuum. Guth's original idea has experienced several complementary modifications [2]. It is the general belief that inflation will finally solve the above mentioned problems of the standard big-bang Cosmology.

The work of Hill, Schramm, and Fry (HSF) [3], renewed the interest in the cosmological significance of domain walls. In the HSF model, the phase transition happens after the time of recombination of matter and radiation. So, domain walls produced during this phase transition are very light and thick, and are assumed to provide the gravitational field necessary for the clustering of dark matter and baryons after the recombination. Those "soft" walls have promising implications to the Universe [4]. Following this line, an interesting solution of thick plane domain wall to the Einstein field equations has been found recently by Goetz [5] and rederived by Mukherjee [6].

Motivated by the inflationary Universe scenario and the interest of the walls on their own right, in this paper we shall present a detailed study of the Goetz solution and pay main attention on its global properties. It is found that for some choice of the free parameter q appearing in the solution, the space-time has a black hole structure but with plane symmetry. It has two asymptotically flat regions separated from the catastrophic ones by event horizons. In each of the two asymptotically flat regions there is a wall. However, a more careful study shows that a Minkowski-like observer will find that the space-time actually has spherical symmetry, and that the walls are bubbles which are collapsing initially and expanding later with a constant acceleration. In this respect, we can see that the thick domain wall shares the same property as those with zero-thickness [7, 8]. Examples of single solutions with several different physical interpretations are not rare in General Relativity [9]. It should be noted that domain walls with a causal lattice structure similar to the Reissner-Nordstrom and Kerr black holes have been found in Supergravity [10], and some interesting features have been obtained [10, 11].

The Goetz solution can be written as [5, 6]

$$ds^2 = e^{-\Omega}(dt^2 - dz^2) - e^{-h}(dx^2 + dy^2), \quad (1)$$

with

$$\Omega = 2q \ln(\cosh(pz)), \quad h = 2q \ln(\cosh(pz)) - 2kt, \quad (2)$$

where $p \equiv k/q$, and k and q are arbitrary constants subject to the conditions $k > 0$, and $0 < q < 1$. The coordinates take the range $-\infty < t, z, x, y < +\infty$. From Eq.(1) we see that Goetz wall is plane symmetric with the Killing vectors, ∂_x, ∂_y , and $y\partial_x - x\partial_y$. The corresponding energy-momentum tensor (EMT) is given by [12] $T_{\mu\nu} = \rho(g_{\mu\nu} + \xi_\mu \xi_\nu) + \nu \xi_\mu \xi_\nu$, where ρ denotes the energy density of the wall, ν the pressure in the direction perpendicular to the wall, and are given by

$$\rho = - \left(\frac{q+2}{3q} \right) \nu = k^2 \left(\frac{q+2}{q} \right) \{\cosh(pz)\}^{-2(1-q)}. \quad (3)$$

The unity vector ξ_μ is the normal to the wall and given by $\xi_\mu = e^{-\Omega/2} \delta_\mu^z$. As shown in [5, 6], the above solution corresponds to a Higgs scalar field with a kinklike shape $\phi (= \text{Arctan}[\sinh(pz)])$, self-interacting through the potential $V(\phi) = \{\cos^2 \phi\}^{(1-q)}$. An interesting feature of this solution is the existence of coordinate singularities in each of the three spatial directions. The ones in the x and y directions are obvious, since the hypersurfaces $z = \text{Const.}$ are the $(2+1)$ -dimensional de Sitter spaces. Thus, these coordinate singularities are the usual de Sitter horizons. The coordinate singularities in the z direction can be seen by computing the proper distance between $z = 0$, the center of the wall, and $|z| = \infty$, which is found finite [5]. Consequently, the hypersurfaces

$|z| = \infty$ represent coordinate singularities, too. Therefore, in general an extension in each of these three directions is needed. The extension in the x and y directions is simple and similar to its four-dimensional analog of the de Sitter space-time given in [13]. Thus, in the following we shall restrict ourselves only to the extension in the z direction.

It will be useful to consider first the hypersurfaces $|z| = \infty$ in more details, specially the timelike geodesics perpendicular to the wall. From the first integral, it is found [5, 6] that the time-like geodesic equations yield $dt/d\tau = E \cosh^{2q}(pz)$, $dz/d\tau = \pm \cosh^q(pz) \{E^2 \cosh^{2q}(pz) - 1\}^{1/2}$, $dx/d\tau = 0$, and $dy/d\tau = 0$, where E is the energy of the test particle, and τ the proper time. Perpendicular to the timelike vector $\lambda_{(0)}^\mu (\equiv dx^\mu/d\tau)$, we have other three linearly independent spacelike vectors $\lambda_{(a)}^\mu (a = 1, 2, 3)$ defined by $\lambda_{(1)}^\mu = (dz/d\tau)\delta_t^\mu + (dt/d\tau)\delta_z^\mu$, $\lambda_{(2)}^\mu = e^{h/2}\delta_x^\mu$, and $\lambda_{(3)}^\mu = e^{h/2}\delta_y^\mu$, where h is given by Eq.(2). One can show that such defined four unity vectors form an orthogonal tetrad and have the properties $\lambda_{(0)}^\mu{}_{;\nu} \lambda_{(0)}^\nu = 0 = \lambda_{(a)}^\mu{}_{;\nu} \lambda_{(0)}^\nu$. Thus, the three vectors $\lambda_{(a)}^\mu$ are parallel transported along the timelike geodesics, and together with $\lambda_{(0)}^\mu$ form a freely-falling frame. Computing the Riemann tensor in this frame, we find that it has only four independent components, one of which is given by

$$R_{\mu\nu\sigma\delta} \lambda_{(0)}^\mu \lambda_{(2)}^\nu \lambda_{(0)}^\sigma \lambda_{(2)}^\delta = \frac{k^2}{q} (\cosh(pz))^{2(q-1)} \{E^2(1-q) \cosh^{2q}(pz) - 1\}.$$

Clearly, as $|z| \rightarrow \infty$, this component becomes unbounded for $q > 1/2$. From the geodesic equation we get that as $|z| \rightarrow \infty$, $e^{2pqz} \sim (\tau_\infty - \tau)^{-1}$. Thus the freely-falling observer experiences a tidal force $\sim (\tau_\infty - \tau)^{-2(q-1/2)/q}$, where τ_∞ is the observer proper time needed to reach $|z| = \infty$, which is finite. From the above expression we can see that the tidal forces experienced by the freely-falling test particles are infinitely large as the hypersurfaces $|z| = \infty$ are approaching. This means that these surfaces are real space-time singularities for $q > 1/2$, instead of horizons [5]. Since in the present case all the scalar invariants are finite, we conclude that these singularities are intermediate (or non-scalar) singularities [14, 15]. On the other hand, the surface energy density of the wall per surface element is given by [5, 6]

$$\sigma = \int_{-\infty}^{\infty} \rho(z) e^{-\Omega/2} dz = \sqrt{\pi} k (2 + q) \frac{\Gamma(1 - q/2)}{\Gamma(3/2 - q/2)},$$

where $\Gamma(x)$ is the standard Gamma function. From this expression we can see that σ is a monotonically increasing function of q . Note that the constant k has the meaning of the energy scale. So, we have that *when the surface energy density of the wall per surface element is greater than σ_c ($\equiv \sigma|_{q=1/2}$), the space-time will be closed at a finite proper distance by space-time singularities*. As King [16] suggested, the non-scalar singularities might not be stable against perturbations and give rise to scalar ones. The following analysis is in favor to King's conjecture, and will show that they are indeed turned into scalar

ones when null fluids are present. In this vein, following [17] we make the substitution

$$\{\Omega, h\} \rightarrow \{\Omega + a(u) + b(v), h\},$$

in the metric coefficients of Eq.(1), where $a(u)$ and $b(v)$ are arbitrary functions of their indicated arguments, with $u \equiv (t + z)/\sqrt{2}$, and $v \equiv (t - z)/\sqrt{2}$. Then, corresponding to the new solution, the EMT is given by

$$T_\nu^\mu = \rho_1 l^\mu l_\nu + \rho_2 n^\mu n_\nu + \tilde{\rho}(\delta_\nu^\mu + \tilde{\xi}^\mu \tilde{\xi}_\nu) + \tilde{\nu} \tilde{\xi}^\mu \tilde{\xi}_\nu, \quad (4)$$

where

$$\begin{aligned} \rho_1 &= -\sqrt{2}k(1 - \tanh(pz))a'(u), & \rho_2 &= -\sqrt{2}k(1 + \tanh(pz))b'(v), \\ \tilde{\rho} &= e^{a(u)+b(v)}\rho, & \tilde{\nu} &= e^{a(u)+b(v)}\nu, \end{aligned} \quad (5)$$

$\tilde{\xi}_\mu = e^{-(a+b)/2}\xi_\mu$, and l_μ and n_μ are two null vectors defined, respectively, by $l_\mu \equiv \partial u / \partial x^\mu = (\delta_\mu^t + \delta_\mu^z)/\sqrt{2}$, $n_\mu \equiv \partial v / \partial x^\mu = (\delta_\mu^t - \delta_\mu^z)/\sqrt{2}$. The function ρ_1 represents the energy density of the null fluid moving along the $v = \text{Const.}$ hypersurfaces, and ρ_2 represents the energy density of the null fluid moving along the $u = \text{Const.}$ hypersurfaces. To have $\rho_{1,2}$ positive, in the following we shall assume that $a'(u), b'(v) < 0$. Combining this assumption with Eq.(5) we can see that, because of the backreaction of the null fluids, the energy density and pressure of the wall become time-dependent, and are always decreasing as the time develops. On the other hand, Eq.(5) also shows that

ρ_1 is vanishing exponentially as one leaves from the center of the wall to the positive z direction, while ρ_2 is vanishing as one leaves from the center to the negative z direction. Thus, the new solution represents a domain wall emitting massless particles.

Corresponding to the new solution, the Kretschmann scalar can be written as

$$\mathcal{R} \equiv R^{\mu\nu\lambda\delta} R_{\mu\nu\lambda\delta} = e^{2(a+b)} \{ \mathcal{R}_0 + 4e^{2\Omega} (\Phi_{00}^{(0)} \rho_1 + \Phi_{22}^{(0)} \rho_2) + 2e^{2\Omega} \rho_1 \rho_2 \}, \quad (7)$$

where $\mathcal{R}_0$ is the Kretschmann scalar of the background $\{\Omega, h\}$, and now is given by $\mathcal{R}_0 = 12k^4(1 + q^2)q^{-2}(\cosh(pz))^{4(q-1)}$. $\Phi_{00}^{(0)}$ and $\Phi_{22}^{(0)}$ are the components of the traceless Ricci tensor [18] and now given by $\Phi_{00}^{(0)} = \Phi_{22}^{(0)} = k^2(1 - q)/(2q \cosh^2(pz))$. $\rho_{1,2}$ and Ω are given, respectively, by Eqs.(5) and (2). The first term at the right-hand side of Eq.(7) represents the backreaction of the null fluids to the background. The second represents the interaction of the null fluids with the matter components $\Phi_{00}^{(0)}$ and $\Phi_{22}^{(0)}$, which now is proportional to $\cosh^{2(2q-1)}(pz)$. Thus, for $q > 1/2$ this term will become unbounded as $|z| \rightarrow \infty$. That is, because of the interaction of the null fluids with the background, the intermediate singularities originally appearing at $|z| = \infty$ are now turned into scalar ones. The last term represents the interaction between the two null fluids, which now is also proportional to $\cosh^{2(2q-1)}(pz)$. Thus, the interaction between these two fluids also turns

those intermediate singularities into scalar ones.

Note that at the level of the Higgs scalar field, the potential corresponding to the new solution becomes $\tilde{V} = e^{a+b}V$. At the field level the new solution is obtained by formally introducing a nontrivial dependence of a coupling “constant” on the coordinates u and v , while the scalar field is kept unchanged. When the functions $a(u) \approx 0$ and $b(v) \approx 0$ the coupling “constant” is effectively unity. Note that the above conclusions regarding to the singular behavior of the space-time at the space-like infinity $|z| = \infty$ hold for any $a(u)$ and $b(v)$.

On the other hand, considering the covariant derivatives of the Riemann tensor in the freely-falling frame, we find

$$\begin{aligned} R_{(i)(j)(k)(l);(k_1)\dots(k_n)} &\rightarrow \text{Exp}\{[2(2q-1) + 2nq]p|z|\} \\ &\rightarrow (\tau_\infty - \tau)^{-[n+2(q-1/2)/q]}, \quad \text{as } |z| \rightarrow \infty, \end{aligned}$$

where indices inside parentheses denote tetrad components. Clearly, for any given q the derivatives upto certain order will become singular on the hypersurfaces $|z| = \infty$. In particular, for $q > \frac{1}{3}$ the first-order derivatives will become unbounded. This indicates the existence of mild singularities on these surfaces even for the solution with $0 < q \leq \frac{1}{2}$. For example, for $q > \frac{1}{3}$ we have that the “difference” of tidal forces become infinitely large but the integral on the surface is still finite. Therefore, one cannot exclude the possibility of

an extension for $0 < q \leq \frac{1}{2}$. The above argument can be further justified by the following consideration. As we shall show below, the Riemann tensor for the extended solution is C^r across $|z| = \infty$ with $r > 0$. According to the classifications given in [14], these hypersurfaces are C^r regular surfaces.

Following [19] (see also [10]), we first make the coordinate transformation

$$u = \alpha^{-1} e^{-k(t+|z|)}, \quad v = -\alpha^{-1} e^{k(t-|z|)}, \quad (8)$$

where $\alpha^2 = 2k^2 4^{-q}$. Then, in terms of u and v the solution of Eqs.(1) and (2) reads

$$ds^2 = [1 + (-\alpha^2 uv)^{1/q}]^{-2q} \{ 2dudv - 2k^2 v^2 (dx^2 + dy^2) \}, \quad (9)$$

where $0 < q \leq 1/2$. From Eq.(8) we can see that the coordinate transformations are restricted to the regions $uv < 0$. To extend the solution into the whole (u, v) -plane, one just simply forgets the way how to get Eq.(9) and lets u and v be any values. Regarding to such an extension, there are two different points of view. The first is due to Cvetič and co-workers [10], namely, considering it as two coordinate transformations, each of which is independently performed in the regions $z \leq 0$ and $z \geq 0$. The resulting space-time of the wall is the gluing of these two extended spaces along the wall. The second is to consider Eq.(8) as one, and take Eq.(9) as the complete extension of the space-time of the wall. Physically, the latter is equivalent to identify the two extended spaces of the former at the same values of the

coordinates. In the following, we shall adopt the second point of view. From Eq.(9) we conclude that in order to extend the solution to all the values of u and v , we must distinguish different cases depending on the solutions of the algebraic equations $(-1)^{1/q} = -1, 1, i$. We shall consider

$$a) q = \frac{2n+1}{2m+1}, \quad b) q = \frac{2n+1}{2m}, \quad c) \text{ the rest, } (n, m = 0, 1, 2, \dots).$$

We shall study the last case first.

Case c). The metric coefficients of Eq.(9) in this case become complex in the regions $uv > 0$, which indicates that Eq.(9) can not be considered as the proper extension beyond the surfaces $uv = 0$, and other possibilities must be considered. One way is to set the conformal factor in Eq.(9) as $[1 + (\alpha^2 uv)^{1/q}]^{-2q}$ in the regions $uv > 0$. One can show that such an extension is C^1 in the sense of [13] and the hypersurfaces $u = 0$ and $v = 0$ are free of any kind of matter. In the regions $uv > 0$, introducing the coordinates t and z via the relations [cf. Eq.(8)]

$$u = \alpha^{-1} e^{-k(t+|z|)}, \quad v = \alpha^{-1} e^{k(t-|z|)},$$

we find that the metric in these regions can be written as

$$ds^2 = \cosh^{-2q}(pz) \left\{ dz^2 - dt^2 - e^{2kt}(dx^2 + dy^2) \right\}, \quad (uv > 0),$$

which is the continuation of the metric (1) across the hypersurfaces $|z| = \infty$, and clearly shows that the coordinate t becomes space-like and z time-like.

Therefore, the hypersurfaces $|z| = \infty$ are acting as Rindler horizons [20]. It should be noted that the extension in this case is not an analytic extension, but it is maximal, in the sense that the extended space-time is geodesically complete [13]. Other possibilities exist, for example, in the regions $uv > 0$, one can replace the conformal factor in Eq.(9) by 1. This extension is C^1 across $|z| = \infty$ and not analytic. The space-time in these regions are flat and the scalar field becomes constant.

Case b). The metric coefficients of Eq.(9) in this case are well defined for all the values of u and v . It can be shown that when $\frac{1}{q}$ is an integer, the extension is the maximal analytic extension, and the extended space-time is geodesically complete. When $\frac{1}{q}$ is not an integer, the extension is only a maximal extension but not analytic. The space-time in the extended regions [cf. Fig.1] is asymptotically flat.

Case a). This is the most interesting case, as after the extension it yields a black hole space-time structure, which is quite similar to that of the Schwarzschild space. The extension in this case is the maximal analytic extension when $\frac{1}{q}$ is an integer, and only a maximal extension otherwise. To show that the space-time indeed has a black hole structure, let us first note that the metric coefficients become singular on the hypersurfaces $uv = \alpha^{-2}$

in this case. On the surface $uv = \alpha^{-2}$ the Kretschmann scalar,

$$\mathcal{R} \equiv R^{\mu\nu\lambda\delta}R_{\mu\nu\lambda\delta} = \frac{48\alpha^4(1+q^2)}{q^2} \frac{(\alpha^2 uv)^{2(1-q)/q}}{[1 - (\alpha^2 uv)^{1/q}]^{4(1-q)}},$$

becomes unbounded. In other words, a space-time singularity appears in the extended regions, II and II' [cf. Fig.1]. From Fig.1 we can see that this singularity is spacelike. Note that in the space-time essentially we have two walls, each of which is located on one of the two branches of the hyperbola $uv = -\alpha^{-2}$. These walls are causally disconnected one from the other and behave like the Rindler particles [20]. The horizons at $|z| = \infty$ (or equivalently $uv = 0$) are event horizons. Thus, it is concluded that the solution given by Eq.(9) in this case represents a black hole but with plane symmetry. The plane is defined by the three Killing vectors ∂_x, ∂_y , and $x\partial_y - y\partial_x$.

However, in [7] (see also [8]) Ipser and Sikivie found that all the plane domain walls with zero-thickness are actually bubbles. The following considerations show explicitly that this is also the case for a thick domain wall. Let us first note that the metric inside the braces of Eq.(9) is flat. As a matter of fact, by performing the following coordinate transformations

$$\begin{aligned} T &= \{(u+v) + k^2 v(x^2 + y^2)\}/\sqrt{2}, \quad Z = \{(u-v) + k^2 v(x^2 + y^2)\}/\sqrt{2}, \\ X &= -\sqrt{2}kvx, \quad Y = -\sqrt{2}kvy, \end{aligned} \tag{10}$$

one can bring this part to the standard Minkowski form. If we further introduce the spherical coordinates $\{R, \theta, \varphi\}$, which are related to the coordinates

$\{T, Z, X, Y\}$ in the usual way, we find that Eq.(9) takes the form

$$ds^2 = \left\{ 1 + \left[\frac{\alpha^2}{2}(R^2 - T^2) \right]^{1/q} \right\}^{-2q} \{dT^2 - dR^2 - R^2(d\theta^2 + \sin^2\theta d\varphi^2)\}, \quad (11)$$

where $R^2 = X^2 + Y^2 + Z^2$. From the above expression we can see that the space-time of the wall has spherical symmetry in the Minkowski coordinates.

On the other hand, from Eq.(10) we have

$$R^2 - T^2 = 2\alpha^{-2}e^{-2k|z|}.$$

Thus, the wall (the center of which is at $z = 0$) essentially is an accelerated bubble with the constant acceleration given by $\alpha/\sqrt{2}$ in the Minkowski coordinates. It starts to collapse at the moment $T = -\infty$ until the moment $T = 0$, where the radius of it is $R_{min.} = \sqrt{2}/\alpha$. Since the acceleration is outward, the wall will start to expand afterwards. Note that the physical radius of the wall is given by $r_{ph} = 2^{-q}R = 2^{-q}(2\alpha^{-2} + T^2)^{1/2}$.

In summary, we have studied the local and global properties of the Goetz thick plane domain wall solution. It has been found that for $1/2 < q < 1$, intermediate singularities appear on the hypersurfaces $|z| = \infty$. When null fluids are present, these singularities become scalar ones. The solution with $0 < q \leq 1/2$ has been extended beyond the horizons $|z| = \infty$. The extended space-time for some choice of the free parameter q has a black hole structure. Moreover, we have shown explicitly that the space-time of the walls in the Minkowski coordinates has spherical symmetry, instead of the plane

symmetry as originally presented. The geometry of the wall is a spherically symmetric closed surface, which is collapsing initially. However, because of its constant outward acceleration, when it collapses to its minimum radius, which is always greater than zero, the wall starts to expand.

Acknowledgments

Part of the work was done when one of the authors (A.W.) was visiting the Department of Applied Mathematics, UNICAMP. He thanks the hospitality of the Department. This work was partially supported by a grant from FAPESP and a grant from CNPq.

References

- [1] A.H. Guth, Phys. Rev. D23, 347 (1981).
- [2] For a recent review on inflationary Universe models, see, for example, E.W. Kolb, Phys. Rep. 227, 5 (1993), and references therein.
- [3] C.T. Hill, D.N. Schramm, and J.N. Fry, Comm. Nucl. Part. Phys. 19, 25 (1989).
- [4] A. Stebbins and M.S. Turner, Astrophys. J. 339, L13 (1989).

- [5] G. Goetz, J. Math. Phys. 31, 2683 (1990).
- [6] M. Mukherjee, Class. Quantum Grav. 10, 131 (1993).
- [7] J. Ipser and P. Sikivie, Phys. Rev. D30, 712 (1984); J. Ipser, 30, 2452 (1984).
- [8] P.S. Letelier and A.Z. Wang, “*Space-Time Defects*,” to appear in J. Math. Phys. (1994).
- [9] W.B. Bonnor, Gen. Rel. Grav. 24, 551 (1992).
- [10] M. Cvetič, S. Griffies, and H.H. Soleng, Phys. Rev. D48, 2613 (1993), S. Griffies, Ph. D. thesis, University of Pennsylvania, Philadelphia, 1993, and references therein.
- [11] G.W. Gibbons, Nucl. Phys. B394, 3 (1993).
- [12] In this paper, the units will be chosen such that $8\pi G = 1 = c$, where G denotes the gravitational constant, and c the speed of light.
- [13] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, England, 1973).
- [14] G.F.R. Ellis and B.G. Schmidt, Gen. Relativ. Gravit. 8, 915 (1977).
- [15] P.S. Letelier, Phys. Rev. Lett. 66, 268 (1991), Class. Quantum Grav. 9, 1707 (1992), Phys. Rev. D47, 1709 (1993).

- [16] A.R. King, Phys. Rev. D11, 763 (1975).
- [17] P.S. Letelier and A.Z. Wang, Phys. Rev. D49, 5105 (1994); Erratum, preprint (1994).
- [18] A. Z. Wang, J. Math. Phys. 33, 1065 (1992).
- [19] A. Z. Wang, Phys. Rev. D48, 2591 (1993).
- [20] W. Rindler, *Essential Relativity* (Springer-Verlag, New York, 1977).

Figure Caption

Fig.1 The projection of the space-time onto the uv -plane. The hypersurfaces $|z| = +\infty$ ($uv = 0$) are event horizons. The center of the walls is at $z = 0$ ($uv = -\alpha^{-2}$). For Case a) where $q = (2n + 1)/(2m + 1)$, the space-time is singular on $uv = \alpha^{-2}$, and consequently regions II and II' represent two catastrophic regions. Regions II, II', III and III' are the four extended regions that are missing in the (t, z) coordinates, where III $\equiv \{uv < -\alpha^{-2}, u > 0\}$, and III' $\equiv \{uv < -\alpha^{-2}, v > 0\}$.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9411020v1>